arXiv:1212.6184v1 [math.RT] 26 Dec 2012

FROM CM-FINITE TO CM-FREE

FAN KONG, PU ZHANG*

ABSTRACT. We prove that the relative Auslander algebra of a CM-finite algebra is CM-free. Thus, the category of the Gorenstein-projective modules over a CM-finite algebra turns out to be the category of the projective modules over a CM-free algebra. We also prove that the Gorenstein defect category of a CM-finite algebra is triangle-equivalent to the singularity category of its relative Auslander algebra. Thus, the Gorenstein defect category of a CM-finite algebra is uniquely determined by the category of its Gorenstein-projective modules.

Key words: Gorenstein-projective module, CM-finite, CM-free, singularity category, Gorenstein defect category.

1. **Introduction**

1.1. Denote by Λ-mod the category of finitely generated left modules over Artin algebra Λ, by $\mathcal{P}(\Lambda)$ the full subcategory of Λ-mod of the projective Λ-modules, and by $\mathcal{GP}(\Lambda)$ the full subcategory of Λ-mod of the Gorenstein-projective Λ-modules (see 2.1 below for the definition). Then $\mathcal{GP}(\Lambda)$ is a Frobenius category with relative projective-injective objects being projective Λ-modules (cf. [AB], [AR], [EJ], [B2]), and hence the stable category $\underline{\mathcal{GP}(\Lambda)}$ of $\mathcal{GP}(\Lambda)$ modulo $\mathcal{P}(\Lambda)$ is a triangulated category ([H1]).

It is clear that $\mathcal{P}(\Lambda) \subseteq \mathcal{GP}(\Lambda)$. If $\mathcal{GP}(\Lambda) = \mathcal{P}(\Lambda)$, then Λ is said to be *CM-free* (cf. [C2]). Clearly, algebras of finite global dimension are CM-free ([EJ, Proposition 10.2.3]). If Λ has only finitely many isoclasses of indecomposable Gorenstein-projective modules, then Λ is said to be *CM-finite* (cf. [B2]). In this case, let $E_1, \cdots, E_n$ be all the pairwise non-isomorphic indecomposable Gorenstein-projective modules, and put $E = \bigoplus\limits_{1\le i\le n} E_i$. We call $(\operatorname{End}_\Lambda E)^{op}$ *the relative Auslander algebra of* Λ, and denote it by $\mathbf{Aus}(\Lambda)$. CM-finite and CM-free algebras have been recently studied by various authors ([B2-3], [BR], [C1-2], [CY], [GZ], [LZ1-2], [R] etc.). In particular, CM finiteness could characterize the simple hypersurface singularity ([BGS], [CPST], [K]).

For a CM-finite algebra Λ, as a consequence of [LZ1, Main Theorem] we know that Λ is Gorenstein (i.e., $\text{p.d.}_\Lambda\Lambda < \infty$ and $\text{p.d.}\Lambda_\Lambda < \infty$) if and only if $\text{gl.dim}\mathbf{Aus}(\Lambda) < \infty$ (see

---

[*] The corresponding author.

2010 Mathematical Subject Classification. 16G10, 16G70, 18E30.

Supported by the NSF of China (11271251), and STCSM (09XD1402500).

also [B3, Corollary 6.8(v)]). In particular, if Λ is a CM-finite Gorenstein algebra, then $\mathbf{Aus}(\Lambda)$ is CM-free. In general we also have

Theorem 1.1. *Let Λ be a CM-finite algebra. Then $\mathbf{Aus}(\Lambda)$ is CM-free.*

Since $\mathrm{Hom}_\Lambda(E, -): \mathcal{GP}(\Lambda) \longrightarrow \mathcal{P}(\mathbf{Aus}(\Lambda))$ is an equivalence, Theorem 1.1 means that the category of the Gorenstein-projective modules over a CM-finite algebra is equivalent to the category of the projective modules over a CM-free algebra.

Put Ω to be the set of CM-finite algebras, and Θ the set of CM-free algebras. Then Theorem 1.1 implies that there is a surjective map $\mathbf{Aus}: \ \Omega \longrightarrow \Theta$, up to Morita equivalence; moreover, the map **Aus** sends CM-finite Gorenstein algebras to algebras of finite global dimension, and sends CM-finite non-Gorenstein algebras to CM-free non-Gorenstein algebras (note that there does not exist a CM-free Gorenstein algebra of infinite global dimension). Graphically, we have



1.2. Let $D^b(\Lambda)$ be the bounded derived category of Λ-mod, and $K^b(\mathcal{P}(\Lambda))$ the bounded homotopy category of $\mathcal{P}(\Lambda)$. The Verdier quotient $D^b(\Lambda)/K^b(\mathcal{P}(\Lambda))$ is called *the singularity category of* Λ and denoted by $D^b_{sg}(\Lambda)$. Then $D^b_{sg}(\Lambda) = 0$ if and only if $\mathrm{gl.dim}\ \Lambda < \infty$. Buchweitz's Theorem ([Buch], Theorem 4.4.1; see also [H2], Theorem 4.6) says that there is an exact embedding $F: \underline{\mathcal{GP}(\Lambda)} \hookrightarrow D^b_{sg}(\Lambda)$ given by $F(G) = G$, where the second G is the corresponding stalk complex at degree 0; and that if Λ is Gorenstein, then F is an equivalence. The converse is also true, i.e., F is an equivalence implies that Λ is Gorenstein (Theorem 6.9(8) in [B1], [BJO], [Zhu]). In general, to measure the difference of $\underline{\mathcal{GP}(\Lambda)}$ and $D^b_{sg}(\Lambda)$, one defines the Gorenstein defect category $D^b_{\text{defect}}(\Lambda) := D^b_{sg}(\Lambda)/\mathrm{Im}F$ (see [BJO]). Then $D^b_{\text{defect}}(\Lambda) = 0$ if and only if Λ is Gorenstein.

Our second main result is as follows.

Theorem 1.2. *Let Λ be a CM-finite algebra. Then there is a triangle-equivalence*

$$D^b_{\text{defect}}(\Lambda) \cong D^b_{sg}(\mathbf{AUS}(\Lambda)).$$

Together with Theorem 1.1 we know that the Gorenstein defect category of a CM-finite algebra can be realized as the singularity category of a CM-free algebra. As a consequence, two CM-finite algebras Λ and Λ' with an equivalence $\mathcal{GP}(\Lambda) \cong \mathcal{GP}(\Lambda')$ as additive categories have the same Gorenstein defect category, i.e., the Gorenstein defect category of a CM-finite algebra is uniquely determined by the category of its Gorenstein-projective modules.

1.3. The paper is organized as follows. In Section 2 we give a result which is needed in the proof of Theorem 1.1. This is a slight generalization of the corresponding result in [Huang], which answers affirmatively a question in [SWSW]. Section 3 is devoted to the proof of Theorem 1.1; and Section 4 is devoted to the proof of Theorem 1.2.

2. **Gorenstein categories**

Throughout the section $\mathcal{A}$ is an abelian category, and $\mathcal{C}$ is a full subcategory of $\mathcal{A}$ which is closed under direct sums. For a complex $X^\bullet$ over $\mathcal{A}$, by $\operatorname{Hom}(\mathcal{C}, X^\bullet)$ is acyclic we mean that $\operatorname{Hom}(C, X^\bullet)$ is acyclic for each $C \in \mathcal{C}$. Similarly for $\operatorname{Hom}(X^\bullet, \mathcal{C})$ being acyclic.

2.1. In [SWSW] the following full subcategory $\mathcal{G}(\mathcal{C})$ of $\mathcal{A}$ is introduced. By definition, an object L of $\mathcal{G}(\mathcal{C})$ is isomorphic to $\operatorname{Im} d^0$, where $X^\bullet = (X^*, d^*)$ is an acyclic complex with $X^i \in \mathcal{C}$ for all $i \in \mathbb{Z}$, such that both $\operatorname{Hom}(\mathcal{C}, X^\bullet)$ and $\operatorname{Hom}(X^\bullet, \mathcal{C})$ are acyclic. In this case $X^\bullet$ is called *a complete $\mathcal{C}$-resolution of L*. Clearly $\mathcal{C} \subseteq \mathcal{G}(\mathcal{C})$.

If Λ is an Artin algebra, $\mathcal{A} = \Lambda$-mod, and $\mathcal{C} = \mathcal{P}(\Lambda)$, then denote $\mathcal{G}(\mathcal{P}(\Lambda))$ by $\mathcal{GP}(\Lambda)$. Call an object in $\mathcal{GP}(\Lambda)$ *a Gorenstein-projective Λ-module*, and a complete $\mathcal{P}(\Lambda)$-resolution *a complete Λ-projective resolution* (cf. [AB], [AR], [EJ]).

S. Sather-Wagstaff, T. Sharif and D. White proved ([SWSW, 4.10]) that if $\mathcal{C}$ is self-orthogonal, then $\mathcal{G}^2(\mathcal{C}) = \mathcal{G}(\mathcal{C})$, where $\mathcal{G}^2(\mathcal{C}) = \mathcal{G}(\mathcal{G}(\mathcal{C}))$. They also asked ([SWSW, 5.8]) whether $\mathcal{G}^2(\mathcal{C}) = \mathcal{G}(\mathcal{C})$ holds for an arbitrary $\mathcal{C}$? Recently Z. Y. Huang ([Huang, Theorem 4.1]) answered this question affirmatively. The following is a slight generalization of Huang's result (by using different method from the one in [Huang]).

Theorem 2.1. *Let $X^\bullet = (X^*, d^*)$ be an acyclic complex with $X^i \in \mathcal{G}(\mathcal{C})$ for all $i \in \mathbb{Z}$. If both $\operatorname{Hom}(\mathcal{C}, X^\bullet)$ and $\operatorname{Hom}(X^\bullet, \mathcal{C})$ are acyclic, then $\operatorname{Im} d^i \in \mathcal{G}(\mathcal{C})$ for all $i \in \mathbb{Z}$.*

2.2. By diagram chasing one can easily verify the following fact. We omit the details.

Lemma 2.2. *Suppose there is a commutative diagram in $\mathcal{A}$ with exact rows δ and η:*

$$\begin{tikzcd}
\delta: & 0 \arrow[r] & X \arrow[r, "f"] \arrow[d, "\alpha"] & Y \arrow[r, "g"] \arrow[d, "\beta"] & Z \arrow[r] \arrow[d, equal] & 0 \\
\eta: & 0 \arrow[r] & X' \arrow[r, "f'"] & Y' \arrow[r, "g'"] & Z \arrow[r] & 0.
\end{tikzcd}$$

Denote by Δ the corresponding short exact sequence $0 \longrightarrow X \xrightarrow{\binom{f}{\alpha}} Y \oplus X' \xrightarrow{(\beta,\, -f')} Y' \longrightarrow 0$. For any $A \in \mathcal{A}$ we have

(*i*) $\operatorname{Hom}(A, \delta)$ *is exact if and only if both* $\operatorname{Hom}(A, \eta)$ *and* $\operatorname{Hom}(A, \Delta)$ *are exact.*

(*ii*) $\operatorname{Hom}(\delta, A)$ *is exact if and only if both* $\operatorname{Hom}(\eta, A)$ *and* $\operatorname{Hom}(\Delta, A)$ *are exact.*

We also need the following technical lemma.

Lemma 2.3. *Let $\delta : 0 \longrightarrow X_1 \xrightarrow{f} U \xrightarrow{g} X_2 \longrightarrow 0$ be an exact sequence with $U \in \mathcal{G}(\mathcal{C})$ such that both* $\operatorname{Hom}(\mathcal{C}, \delta)$ *and* $\operatorname{Hom}(\delta, \mathcal{C})$ *are exact. Let $\eta : 0 \longrightarrow X_1 \xrightarrow{u} Y \xrightarrow{v} V \longrightarrow 0$ be another exact sequence with $V \in \mathcal{G}(\mathcal{C})$ such that* $\operatorname{Hom}(\eta, \mathcal{C})$ *is exact. Then we have the following commutative diagram*

$$\begin{tikzcd}
\delta: & 0 \arrow[r] & X_1 \arrow[r, "f"] \arrow[d, hook, "u"] & U \arrow[r, "g"] \arrow[d] & X_2 \arrow[r] \arrow[d, hook, "u'"] & 0 \\
\delta': & 0 \arrow[r] & Y \arrow[r] & C \arrow[r] & Z \arrow[r] & 0
\end{tikzcd}$$

such that

(*i*) *δ' is exact with $C \in \mathcal{C}$, and both* $\operatorname{Hom}(\mathcal{C}, \delta')$ *and* $\operatorname{Hom}(\delta', \mathcal{C})$ *are exact;*

(*ii*) *u' is a monomorphism with* $\operatorname{Coker} u' \in \mathcal{G}(\mathcal{C})$*, and* $\operatorname{Hom}(u', \mathcal{C})$ *is surjective.*

Proof. Step 1. Since by assumption $U, V \in \mathcal{G}(\mathcal{C})$, by definition there exists two exact sequences $\epsilon_1 : 0 \longrightarrow U \xrightarrow{a} C_1 \xrightarrow{b} L_1 \longrightarrow 0$ and $\epsilon_2 : 0 \longrightarrow V \xrightarrow{c} C_2 \xrightarrow{d} L_2 \longrightarrow 0$ with $C_1, C_2 \in \mathcal{C}, L_1, L_2 \in \mathcal{G}(\mathcal{C})$, such that $\operatorname{Hom}(\mathcal{C}, \epsilon_1)$, $\operatorname{Hom}(\epsilon_1, \mathcal{C})$, $\operatorname{Hom}(\mathcal{C}, \epsilon_2)$ and $\operatorname{Hom}(\epsilon_2, \mathcal{C})$ are exact. Since by assumption $\operatorname{Hom}(\eta, C_1)$ is exact, by considering $af \in \operatorname{Hom}(X_1, C_1)$ we see that there exists morphisms e and e' such that the following diagram commutates:

$$\begin{tikzcd}
0 \arrow[r] & X_1 \arrow[r, "u"] \arrow[d, "f"] & Y \arrow[r, "v"] \arrow[d, dashed, "e"] & V \arrow[r] \arrow[d, dashed, "e'"] & 0 \\
0 \arrow[r] & U \arrow[r, "a"] & C_1 \arrow[r, "b"] & L_1 \arrow[r] & 0.
\end{tikzcd}$$

Put $\alpha = \binom{e}{cv} : Y \longrightarrow C_1 \oplus C_2$, $i = \binom{a}{0} : U \longrightarrow C_1 \oplus C_2$, $\pi = \left(\begin{smallmatrix} b & 0 \\ 0 & 1 \end{smallmatrix}\right) : C_1 \oplus C_2 \longrightarrow L_1 \oplus C_2$, $x = \binom{e'}{c} : V \longrightarrow L_1 \oplus C_2$. Then there exist morphisms β, u', v', y such that the following diagram commutes:

$$\begin{tikzcd}
 & & \eta & \Xi & \eta' & & \\
 & & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d, dashed] & & \\
\delta : & 0 \arrow[r] & X_1 \arrow[r, "f"] \arrow[d, "u"] & U \arrow[r, "g"] \arrow[d, "i"] & X_2 \arrow[r] \arrow[d, dashed, "u'"] & 0 & \\
\delta' : & 0 \arrow[r] & Y \arrow[r, "\alpha"] \arrow[d, "v"] & C_1 \oplus C_2 \arrow[r, "\beta"] \arrow[d, "\pi"] & Z \arrow[r] \arrow[d, dashed, "v'"] & 0 & (*) \\
\Delta : & 0 \arrow[r] & V \arrow[r, "x"] \arrow[d] & L_1 \oplus C_2 \arrow[r, "y"] \arrow[d] & L \arrow[r] \arrow[d] & 0 & \\
 & & 0 & 0 & 0 & &
\end{tikzcd}$$

where $Z = \operatorname{Coker} \alpha$ and $L = \operatorname{Coker} x$. Note that x is monomorphism since c is so. Also, the middle column Ξ is exact since ϵ_1 is exact. Applying Snake Lemma to the left two columns we know that α is monomorphism and that the right column η' is exact. In particular, u' is a monomorphism.

We will prove that the upper two rows of $(*)$ is what we need.

Step 2. Write $y = (\, m, \, -l \,)$. Observe that Δ is exact means that there is the following commutative diagram with exact rows (note that L is the push-out of e' and c):

$$\begin{tikzcd}
\epsilon_2 : & 0 \arrow[r] & V \arrow[r, "c"] \arrow[d, "e'"] & C_2 \arrow[r, "d"] \arrow[d, "l"] & L_2 \arrow[r] \arrow[d, equal] & 0 \\
\zeta : & 0 \arrow[r] & L_1 \arrow[r, "m"] & L \arrow[r, "n"] & L_2 \arrow[r] & 0
\end{tikzcd}$$

Since both $\operatorname{Hom}(\mathcal{C}, \epsilon_2)$ and $\operatorname{Hom}(\epsilon_2, \mathcal{C})$ are exact, it follows from Lemma 2.2 that $\operatorname{Hom}(\mathcal{C}, \zeta)$, $\operatorname{Hom}(\zeta, \mathcal{C})$, $\operatorname{Hom}(\mathcal{C}, \Delta)$ and $\operatorname{Hom}(\Delta, \mathcal{C})$ are all exact.

Since $L_1, L_2 \in \mathcal{G}(\mathcal{C})$, and both $\operatorname{Hom}(\mathcal{C}, \zeta)$ and $\operatorname{Hom}(\zeta, \mathcal{C})$ are exact, it follows from Proposition 4.4 of [SWSW] that $L \in \mathcal{G}(\mathcal{C})$, thus $\operatorname{Coker} u' = L \in \mathcal{G}(\mathcal{C})$.

Step 3. $\forall\ C \in \mathcal{C}$, applying $\operatorname{Hom}(C,-)$ to the right two columns of $(*)$, we get the following commutative diagram with exact rows

$$\begin{tikzcd}
 & \operatorname{Hom}(C,U) \arrow[r] \arrow[d, "\operatorname{Hom}(C,g)"] & \operatorname{Hom}(C,C_1\oplus C_2) \arrow[r, "(C,\pi)"] \arrow[d, "\operatorname{Hom}(C,\beta)"] & \operatorname{Hom}(C,L_1\oplus C_2) \arrow[r] \arrow[d, "\operatorname{Hom}(C,y)"] & 0 \\
0 \arrow[r] & \operatorname{Hom}(C,X_2) \arrow[r] & \operatorname{Hom}(C,Z) \arrow[r] & \operatorname{Hom}(C,L).
\end{tikzcd}$$

(Note that $\operatorname{Hom}(C,\pi)$ is surjective since $\operatorname{Hom}(C,\epsilon_1)$ is exact.) Also $\operatorname{Hom}(C,y)$ and $\operatorname{Hom}(C,g)$ are surjective, by Snake-Lemma we know that $\operatorname{Hom}(C,\beta)$ is surjective. That is, $\operatorname{Hom}(\mathcal{C},\delta')$ is exact.

Step 4. $\forall\ C \in \mathcal{C}$, applying $\operatorname{Hom}(-,C)$ to $(*)$, we have the following commutative diagram with exact rows and columns:

$$\begin{tikzcd}
 & & (\eta',C) & (\Xi,C) & (\eta,C) & \\
 & & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
(\Delta,C): & 0 \arrow[r] & (L,C) \arrow[r, "(y,C)"] \arrow[d, "(v',C)"] & (L_1\oplus C_2,C) \arrow[r, "(x,C)"] \arrow[d, "(\pi,C)"] & (V,C) \arrow[r] \arrow[d, "(v,C)"] & 0 \\
(\delta',C): & 0 \arrow[r] & (Z,C) \arrow[r, "(\beta,C)"] \arrow[d, "(u',C)"] & (C_1\oplus C_2,C) \arrow[r, "(\alpha,C)"] \arrow[d, "(i,C)"] & (Y,C) \arrow[d, "(u,C)"] & \\
(\delta,C): & 0 \arrow[r] & (X_2,C) \arrow[r, "(g,C)"] & (U,C) \arrow[r, "(f,C)"] \arrow[d] & (X_1,C) \arrow[r] \arrow[d] & 0 \\
 & & & 0 & 0 &
\end{tikzcd}$$

(Note that the exactness of $\operatorname{Hom}(\delta,C)$ and $\operatorname{Hom}(\eta,C)$ follows from assumptions; the exactness of $\operatorname{Hom}(\Xi,C)$ follows from the exactness of $\operatorname{Hom}(\epsilon_1,C)$; and the exactness of $\operatorname{Hom}(\Delta,C)$ follows from Step 2). Applying Snake-Lemma to the right two columns we see that both $\operatorname{Hom}(u',C)$ and $\operatorname{Hom}(\alpha,C)$ are surjective. This completes the proof. ■

2.3. **Proof of Theorem 2.1.** Without losing of generality, we only prove $\operatorname{Im} d^0 \in \mathcal{G}(\mathcal{C})$. For each i, decompose d^i as $X^i \overset{a^i}{\twoheadrightarrow} \operatorname{Im} d^i \xhookrightarrow{b^i} X^{i+1}$. We claim that there exists an exact sequence $C^+ : 0 \longrightarrow \operatorname{Im} d^0 \xrightarrow{v^0} C^1 \xrightarrow{\gamma^1} C^2 \xrightarrow{\gamma^2} \cdots$ with $C^i \in \mathcal{C}$ for all $i \geq 1$, such that both $\operatorname{Hom}(C^+,\mathcal{C})$ and $\operatorname{Hom}(\mathcal{C},C^+)$ are exact.

In fact, applying Lemma 2.3 to the exact sequence $0 \longrightarrow \operatorname{Im} d^0 \xrightarrow{b^0} X^1 \xrightarrow{a^1} \operatorname{Im} d^1 \longrightarrow 0$ and the exact sequence $0 \longrightarrow \operatorname{Im} d^0 \xrightarrow{=} \operatorname{Im} d^0 \longrightarrow 0 \longrightarrow 0$, we get the following commutative diagram with exact rows

$$\begin{tikzcd}
 & 0 \arrow[r] & \operatorname{Im} d^0 \arrow[r, "b^0"] \arrow[d, equal] & X^1 \arrow[r, "a^1"] \arrow[d] & \operatorname{Im} d^1 \arrow[r] \arrow[d, hook, "w^1"] & 0 \\
\xi_1 : & 0 \arrow[r] & \operatorname{Im} d^0 \arrow[r, "v^0"] & C^1 \arrow[r, "u^1"] & L^1 \arrow[r] & 0
\end{tikzcd}$$

such that $C^1 \in \mathcal{C}$, w^1 is a monomorphism with $\operatorname{Coker} w^1 \in \mathcal{G}(\mathcal{C})$, and that $\operatorname{Hom}(\mathcal{C}, \xi_1)$ and $\operatorname{Hom}(\xi_1, \mathcal{C})$ are exact, and $\operatorname{Hom}(w^1, \mathcal{C})$ is surjective. Now applying Lemma 2.3 to the exact sequence $0 \longrightarrow \operatorname{Im} d^1 \xrightarrow{b^1} X^2 \xrightarrow{a^2} \operatorname{Im} d^2 \longrightarrow 0$ and the exact sequence $0 \longrightarrow \operatorname{Im} d^1 \xrightarrow{w^1} L^1 \longrightarrow \operatorname{Coker} w^1 \longrightarrow 0$, we get an exact sequence $\xi_2 : 0 \longrightarrow L_1 \xrightarrow{v^1} C_2 \xrightarrow{u^2} L_3 \longrightarrow 0$ and monomorphism $w^2 : \operatorname{Im} d^2 \hookrightarrow L^2$. Continuing this process and putting $\xi_1, \xi_2, \cdots$ together we finally obtain an exact sequence C^+ with $\gamma^i = v^i u^i, \ i \geq 1$, such that $\operatorname{Hom}(C^+, \mathcal{C})$ and $\operatorname{Hom}(\mathcal{C}, C^+)$ are exact (since $\operatorname{Hom}(\xi_i, \mathcal{C})$ and $\operatorname{Hom}(\mathcal{C}, \xi_i)$ are exact for all $i \geq 1$).

Dually, there exists an exact sequence $C^- : \cdots \longrightarrow C^{-1} \longrightarrow C^0 \longrightarrow \operatorname{Im} d^0 \longrightarrow 0$ with $C^i \in \mathcal{C}$ for $i \leq 0$ such that both $\operatorname{Hom}(C^-, \mathcal{C})$ and $\operatorname{Hom}(\mathcal{C}, C^-)$ are exact (this can be done shortly by applying the above claim to the opposite category $\mathcal{A}^{\mathrm{op}}$). Now putting C^- and C^+ together, we obtain a complete $\mathcal{C}$-resolution of $\operatorname{Im} d^0$, and this proves $\operatorname{Im} d^0 \in \mathcal{G}(\mathcal{C})$. ■

3. **Proof of Theorem 1.1**

3.1. Applying Theorem 2.1 to the case $\mathcal{A} = \Lambda$-mod and $\mathcal{C} = \mathcal{P}(\Lambda)$, where Λ is an Artin algebra, we get the following

Proposition 3.1. *Let* $X^\bullet = (X^*, d^*)$ *be an acyclic complex with* $X^i \in \mathcal{GP}(\Lambda)$ *for all* i. *If* $\operatorname{Hom}_\Lambda(X^\bullet, \Lambda)$ *is still acyclic, then* $\operatorname{Im} d^i \in \mathcal{GP}(\Lambda)$ *for all* i.

3.2. Let E be a Gorenstein-projective generator, i.e., E is a Gorenstein-projective Λ-module such that $\Lambda \in \operatorname{add} E$. For convenience, denote $\mathcal{G}(\operatorname{add} E)$ by $\mathcal{GP}(E)$, i.e.,

$$\begin{aligned}\mathcal{GP}(E) = \{X \in \Lambda\text{-}mod \ | \ &\exists \text{ an acyclic complex } X^\bullet = (X^*, d^*) \text{ with all } X^i \in \operatorname{add} E \\ &\text{such that } \operatorname{Hom}(X^\bullet, E) \text{ and } \operatorname{Hom}(E, X^\bullet) \text{ are again acyclic, and } X \cong \operatorname{Im} d^0\}.\end{aligned}$$

By Proposition 3.1 we have

Corollary 3.2. *Let* E *be a Gorenstein-projective generator. Then* $\mathcal{GP}(E) \subseteq \mathcal{GP}(\Lambda)$.

3.3. Recall the Yoneda philosophy. Given a Λ-module E, then the functor $\operatorname{Hom}_\Lambda(E, -)$: Λ-mod $\longrightarrow \Gamma$-mod induces an equivalence between $\operatorname{add} E$ and $\mathcal{P}(\Gamma)$, where $\Gamma = (\operatorname{End}_\Lambda E)^{op}$. Also, for each $E' \in \operatorname{add} E$ and each $X \in \Lambda$-mod there is an isomorphism

$$\operatorname{Hom}_\Lambda(E', X) \longrightarrow \operatorname{Hom}_\Gamma(\operatorname{Hom}_A(E, E'), \operatorname{Hom}_A(E, X))$$

given by $f \mapsto \operatorname{Hom}_\Lambda(E, f), \ \forall \ f \in \operatorname{Hom}_\Lambda(E', X)$ (cf. [ARS], p.33).

Now, if E is a generator (i.e., $\Lambda \in \operatorname{add} E$), then we can say more.

Lemma 3.3. *Let E be a generator of Λ-mod, and $\Gamma = (\operatorname{End} E)^{op}$. Then the functor* $\operatorname{Hom}_\Lambda(E, -) : \Lambda\text{-mod} \longrightarrow \Gamma\text{-mod}$ *is fully faithful.*

Proof. Since E is a generator, for any Λ-module X there is a surjective map $E^m \twoheadrightarrow X$ for some positive integer m. This implies that $\operatorname{Hom}_\Lambda(E, -)$ is faithful. Let $X, Y \in \Lambda$-mod, $f : \operatorname{Hom}_\Lambda(E, X) \longrightarrow \operatorname{Hom}_\Lambda(E, Y)$ be a Γ-map. By taking right $\operatorname{add} E$-approximations, we get exact sequences $T_1 \xrightarrow{u} T_0 \xrightarrow{\pi} X \longrightarrow 0$ and $T'_1 \xrightarrow{u'} T'_0 \xrightarrow{\pi'} Y \longrightarrow 0$ with $T_0, T_1, T'_0, T'_1 \in \operatorname{add} E$ (note that since E is a generator, it follows that π and π' are surjective). Applying $\operatorname{Hom}_\Lambda(E, -)$ we have the following diagram with exact rows

$$\begin{tikzcd}
\operatorname{Hom}(E, T_1) \arrow[r, "\operatorname{Hom}(E,u)"] \arrow[d, dashed, "f_1"] & \operatorname{Hom}(E, T_0) \arrow[r, "\operatorname{Hom}(E,\pi)"] \arrow[d, dashed, "f_0"] & \operatorname{Hom}(E, X) \arrow[r] \arrow[d, "f"] & 0 \\
\operatorname{Hom}(E, T'_1) \arrow[r, "\operatorname{Hom}(E,u')"] & \operatorname{Hom}(E, T'_0) \arrow[r, "\operatorname{Hom}(E,\pi')"] & \operatorname{Hom}(E, Y) \arrow[r] & 0.
\end{tikzcd}$$

Since the two rows are respectively parts of projective resolutions of $\operatorname{Hom}(E, X)$ and $\operatorname{Hom}(E, Y)$, f induces f_1 and f_0 such that the above diagram commutes. Thus $f_i = \operatorname{Hom}(E, f'_i)$ for some $f'_i \in \operatorname{Hom}_\Lambda(T_i, T'_i)$, $i = 0, 1$. So we get the following diagram

$$\begin{tikzcd}
T_1 \arrow[r, "u"] \arrow[d, "f'_1"] & T_0 \arrow[r, "\pi"] \arrow[d, "f'_0"] & X \arrow[r] \arrow[d, dashed, "f'"] & 0 \\
T'_1 \arrow[r, "u'"] & T'_0 \arrow[r, "\pi'"] & Y \arrow[r] & 0
\end{tikzcd}$$

with commutative left square. So there exists $f' \in \operatorname{Hom}_\Lambda(X, Y)$ such that the above diagram commutes. Thus $f \operatorname{Hom}(E, \pi) = \operatorname{Hom}(E, f') \operatorname{Hom}(E, \pi)$ and hence $f = \operatorname{Hom}(E, f')$. This proves that $\operatorname{Hom}(E, -)$ is full. ■

3.4. The following result shows that, after taking the opposite algebra of endomorphism algebra of a Gorenstein-projective generator, the category of the Gorenstein-projective modules can not be "enlarged". It is a main step for proving Theorem 1.1.

Theorem 3.4. *Let E be a Gorenstein-projective generator, and $\Gamma = (\operatorname{End}_\Lambda E)^{op}$. Then* $\operatorname{Hom}_\Lambda(E, -)$ *induces an equivalence between $\mathcal{GP}(E)$ and $\mathcal{GP}(\Gamma)$. In particular, $\mathcal{GP}(\Gamma)$ is equivalent to a full subcategory of $\mathcal{GP}(\Lambda)$.*

Proof. Let $L \in \mathcal{GP}(E)$. By definition there exists an acyclic complex $X^\bullet = (X^*, d^*)$ with all $X^i \in \operatorname{add} E$ such that $\operatorname{Hom}_\Lambda(E, X^\bullet)$ and $\operatorname{Hom}_\Lambda(X^\bullet, E)$ are acyclic, and $L \cong \operatorname{Im} d^0$. Since $\operatorname{Hom}_\Lambda(X^\bullet, E) \cong \operatorname{Hom}_\Gamma(\operatorname{Hom}_\Lambda(E, X^\bullet), \Gamma)$, it follows that $\operatorname{Hom}_\Gamma(\operatorname{Hom}_\Lambda(E, X^\bullet), \Gamma)$ is acyclic. Thus $\operatorname{Hom}_\Lambda(E, X^\bullet)$ is a complete Γ-projective resolution of $\operatorname{Im}(\operatorname{Hom}_\Lambda(E, d^0))$. However, Since $\operatorname{Hom}_\Lambda(E, X^\bullet)$ is acyclic and E is a generator, it follows that $\operatorname{Im}(\operatorname{Hom}_\Lambda(E, d^0))$ $= \operatorname{Hom}_\Lambda(E, \operatorname{Im} d^0)$. That is, $\operatorname{Hom}_\Lambda(E, L) \in \mathcal{GP}(\Gamma)$.

We claim that $\operatorname{Hom}_\Lambda(E,-) : \mathcal{GP}(E) \longrightarrow \mathcal{GP}(\Gamma)$ is dense. Suppose $Y \in \mathcal{GP}(\Gamma)$. Then there is a complete projective Γ-resolution $P^\bullet : \cdots \longrightarrow P^{-1} \xrightarrow{d^{-1}} P^0 \xrightarrow{d^0} P^1 \xrightarrow{d^1} \cdots$ such that $Y \cong \operatorname{Im} d^0$. Then there exists a complex $E^\bullet : \cdots \longrightarrow E^{-1} \xrightarrow{e^{-1}} E^0 \xrightarrow{e^0} E^1 \xrightarrow{e^1} \cdots$ with all $E^i \in \operatorname{add} E$ such that $\operatorname{Hom}_\Lambda(E, E^\bullet) = P^\bullet$. Since E is a generator, for each i there is a surjective Λ-map $\pi : E' \twoheadrightarrow \operatorname{Ker} e^{i+1}$ with $E' \in \operatorname{add} E$; while $\operatorname{Hom}_\Lambda(E', E^\bullet)$ is acyclic, so there is $g \in \operatorname{Hom}(E', E^i)$ such that $\pi = e^i g$. Decomposing e^i as the composition $E^i \twoheadrightarrow \operatorname{Im} e^i \hookrightarrow \operatorname{Ker} e^{i+1}$, we see that the embedding $\operatorname{Im} e^i \hookrightarrow \operatorname{Ker} e^{i+1}$ is surjective. This proves that $E^\bullet$ is also acyclic. Since $\operatorname{Hom}_\Gamma(P^\bullet, \Gamma) \cong \operatorname{Hom}_\Lambda(E^\bullet, E)$ and $\operatorname{Hom}_\Gamma(\Gamma, P^\bullet) \cong \operatorname{Hom}_\Lambda(E, E^\bullet)$, it follows that $\operatorname{Hom}_\Lambda(E^\bullet, E)$ and $\operatorname{Hom}_\Lambda(E, E^\bullet)$ are acyclic, and hence $\operatorname{Im} e^i \in \mathcal{GP}(E)$ for all i. Since $\operatorname{Hom}_\Lambda(E, E^\bullet)$ is acyclic, $\operatorname{Im} d^i \cong \operatorname{Im}(\operatorname{Hom}_\Lambda(E, e^i)) = \operatorname{Hom}_\Lambda(E, \operatorname{Im} e^i)$. Thus $Y \cong \operatorname{Im} d^0 \in \operatorname{Hom}_\Lambda(E, \mathcal{GP}(E))$. This proves the claim.

By Lemma 3.3 $\operatorname{Hom}_\Lambda(E,-) : \Lambda\text{-mod} \longrightarrow \Gamma\text{-mod}$ is fully faithful. Thus $\operatorname{Hom}_\Lambda(E,-) : \mathcal{GP}(E) \longrightarrow \mathcal{GP}(\Gamma)$ is an equivalence of categories. Since $\mathcal{GP}(E) \subseteq \mathcal{GP}(\Lambda)$ by Corollary 3.2, we see that $\mathcal{GP}(\Gamma)$ is equivalent to a full subcategory of $\mathcal{GP}(\Lambda)$. ■

Corollary 3.5. *Let Λ be a CM-finite Artin algebra, E a Gorenstein-projective generator. Then $\Gamma = (\operatorname{End} E)^{op}$ is also CM-finite.*

3.5. Now we can prove Theorem 1.1. Instead we prove the following version of Theorem 1.1, which is more precise. We stress that the assertion (ii) is well-known.

Theorem 3.6. *Let Λ be a CM-finite algebra, $E_1, \ldots, E_n$ are all the pairwise non-isomorphic indecomposable Gorenstein-projective Λ-modules. Put $E = \bigoplus_{1\le i\le n} E_i$ and $\mathbf{Aus}(\Lambda) = (\operatorname{End} E)^{op}$. Then $\mathbf{Aus}(\Lambda)$ is CM-free.*

More precisely, put Ω to be the set of CM-finite algebras, and Θ the set of CM-free algebras. Then

(i) The map $\mathbf{Aus} :\ \Omega \longrightarrow \Theta$ is surjective, up to Morita equivalence.

(ii) The map $\mathbf{Aus}$ sends CM-finite Gorenstein algebras to algebras of finite global dimension.

(iii) The map $\mathbf{Aus}$ sends CM-finite non-Gorenstein algebras to CM-free non-Gorenstein algebras. In particular, there does not exist a CM-free Gorenstein algebra of infinite global dimension.

Proof. Write $\Gamma = \mathbf{Aus}(\Lambda)$. By definition we have $\mathcal{GP}(\Lambda) = \operatorname{add} E \subseteq \mathcal{GP}(E)$. By Corollary 3.2 $\mathcal{GP}(E) \subseteq \mathcal{GP}(\Lambda)$. Thus $\mathcal{GP}(E) = \operatorname{add} E$. By Theorem 3.4 we have

$$\mathcal{GP}(\Gamma) = \operatorname{Hom}_\Lambda(E,\ \mathcal{GP}(E)) = \operatorname{Hom}_\Lambda(E,\ \operatorname{add} E) = \mathcal{P}(\Gamma).$$

Thus Γ is CM-free. Note that the map $\mathbf{Aus}$ sends a CM-free algebra to itself up to Morita equivalence, and hence the assertion (i) follows.

Recall a well-known fact: for a CM-finite algebra Λ, Λ is Gorenstein if and only if Γ is of finite global dimension (this is a consequence of Main Theorem in [LZ1]. See also [B3, Corollary 6.8(v)]). From this the assertion (ii) follows.

By the fact stated above, **Aus** sends CM-finite non-Gorenstein algebra Λ to a CM-free algebra Γ of infinite global dimension. Note that Γ can not be Gorenstein: otherwise, by (ii) we see that Γ is of finite global dimension, since Γ is Morita equivalent to $\mathbf{Aus}(\Gamma)$. This proves (iii). ■

3.6. Let E be a Gorenstein-projective generator of Λ-mod. By Corollary 3.2 we have $\mathcal{GP}(E) \subseteq \mathcal{GP}(\Lambda)$; and by Theorem 3.4 if $\mathcal{GP}(E) = \mathcal{GP}(\Lambda)$ then $\operatorname{Hom}_\Lambda(E, -)$ induces an equivalence between $\mathcal{GP}(E)$ and $\mathcal{GP}(\Gamma)$, where $\Gamma = (\operatorname{End}_\Lambda E)^{op}$. So, we further discuss the condition $\mathcal{GP}(E) = \mathcal{GP}(\Lambda)$, in the rest of this section.

Denote by $\langle E\rangle$ the set of morphisms in Λ-mod factoring through $\operatorname{add} E$, and by $\mathcal{GP}(E)/\langle E\rangle$ the corresponding factor category. Note that the usual stable category $\underline{\mathcal{GP}(\Lambda)} = \mathcal{GP}(\Lambda)/\langle\Lambda\rangle$ is a triangulated category. The following fact shows that one may construct different triangulated structures, starting from $\mathcal{GP}(\Lambda)$.

Proposition 3.7. *Let E be a Gorenstein-projective generator of Λ-mod such that $\mathcal{GP}(E) = \mathcal{GP}(\Lambda)$. Then $\mathcal{GP}(\Lambda)/\langle E\rangle$ is a triangulated category.*

Proof. Put $\Gamma = (\operatorname{End} E)^{op}$. By Theorem 3.4 we have equivalence $\mathcal{GP}(E) \cong \mathcal{GP}(\Gamma)$ with restriction $\operatorname{add} E \cong \mathcal{P}(\Gamma)$. Thus we get an equivalence $\mathcal{GP}(E)/\langle E\rangle \cong \mathcal{GP}(\Gamma)/\langle\Gamma\rangle = \underline{\mathcal{GP}(\Gamma)}$, from which and the assumption the assertion follows. ■

In order to describe when $\mathcal{GP}(E) = \mathcal{GP}(\Lambda)$, we introduce the following

Definition 3.8. *Suppose E is a Gorenstein-projective generator. If the following two conditions are satisfied, then we call E a center of $\mathcal{GP}(\Lambda)$:*

(i) For each $X \in \mathcal{GP}(\Lambda)$, there exists an exact sequence $0 \longrightarrow X \xrightarrow{f} E' \xrightarrow{g} Y \longrightarrow 0$ such that f is a left $\operatorname{add} E$-approximation and g is a right $\operatorname{add} E$-approximation.

(ii) For each $X \in \mathcal{GP}(\Lambda)$, there exists an exact sequence $0 \longrightarrow Y \xrightarrow{f} E' \xrightarrow{g} X \longrightarrow 0$ such that f is a left $\operatorname{add} E$-approximation and g is a right $\operatorname{add} E$-approximation.

It is clear that in (ii) of Definition 3.8 we have $Y \in \mathcal{GP}(\Lambda)$; we stress that in (i) of Definition 3.8 we also have $Y \in \mathcal{GP}(\Lambda)$.

We have the following description.

Proposition 3.9. *Suppose E is a Gorenstein-projective generator. Then $\mathcal{GP}(E) = \mathcal{GP}(\Lambda)$ if and only if E is a center of $\mathcal{GP}(\Lambda)$.*

Proof. "$\Longrightarrow$": For each $X \in \mathcal{GP}(\Lambda) = \mathcal{GP}(E)$, there exists an acyclic complex $X^\bullet = (X^*, d^*)$ with all $X^i \in \operatorname{add} E$, such that $\operatorname{Hom}(X^\bullet, E)$ and $\operatorname{Hom}(E, X^\bullet)$ are again acyclic, and $X = \operatorname{Im} d^0$. Thus the exact sequences $0 \longrightarrow X \longrightarrow X^1 \longrightarrow \operatorname{Im} d^1 \longrightarrow 0$ and $0 \longrightarrow \operatorname{Im} d^{-1} \longrightarrow X^0 \longrightarrow X \longrightarrow 0$ satisfy the conditions in Definition 3.8.

"$\Longleftarrow$": Given $L = L^0 \in \mathcal{GP}(\Lambda)$, by induction on i, there exists exact sequences $0 \longrightarrow L^i \xrightarrow{v^i} X^{i+1} \xrightarrow{u^{i+1}} L^{i+1} \longrightarrow 0$ and $0 \longrightarrow L^{-i-1} \xrightarrow{v^{-i-1}} X^{-i} \xrightarrow{u^{-i}} L^{-i} \longrightarrow 0$ with $L^i, L^{i+1}, L^{-i}, L^{-i-1} \in \mathcal{GP}(\Lambda)$, such that v^i and v^{-i-1} are left $\operatorname{add} E$-approximations, and u^{i+1} and u^{-i} are right $\operatorname{add} E$-approximations. Combining them we obtain a complete $\operatorname{add} E$-resolution of L. Together with Corollary 3.2 this proves $\mathcal{GP}(E) = \mathcal{GP}(\Lambda)$. ■

3.7. The above discussion is of particular interest in the case that Λ is a self-injective algebra. In this case any Λ-module is Gorenstein-projective.

Corollary 3.10. *Let Λ be a self-injective algebra, E a generator. Then $\mathcal{GP}(E) = \mathcal{GP}(\Lambda)$ if and only if* $\operatorname{DTr} E \oplus \operatorname{TrD} E \in \operatorname{add} E$.

Proof. By Proposition 3.9 we just prove that E is a center of $\mathcal{GP}(\Lambda)$ if and only if $\operatorname{DTr} E \oplus \operatorname{TrD} E \in \operatorname{add} E$.

Suppose $\operatorname{DTr} E \oplus \operatorname{TrD} E \in \operatorname{add} E$. Given $X \in \mathcal{GP}(\Lambda) = \Lambda\text{-mod}$, let $0 \longrightarrow X \xrightarrow{f} E' \xrightarrow{g} Y \longrightarrow 0$ be an exact sequence such that f is a left $\operatorname{add} E$-approximation of X. Then $\operatorname{Hom}_\Lambda(f, \operatorname{DTr} E)$ is epic. By Theorem 4.1 in [ARS] we see that $\operatorname{Hom}_\Lambda(E, g)$ is epic, i.e., g is a right $\operatorname{add} E$-approximation of Y. So E satisfies the condition (i) of Definition 3.8. Dually, E satisfies the condition (ii), and hence E is a center.

Conversely, suppose E is a center. Then there is an exact sequence $0 \longrightarrow \operatorname{DTr} E \xrightarrow{i} E' \xrightarrow{\pi} L \longrightarrow 0$ such that i is a left $\operatorname{add} E$-approximation of $\operatorname{DTr} E$ and π is a right $\operatorname{add} E$-approximation of L. However by Theorem 4.1 in [ARS] we know $\operatorname{Hom}_\Lambda(i, \operatorname{DTr} E)$ is epic since $\operatorname{Hom}_\Lambda(E, \pi)$ is epic. Hence i splits and $\operatorname{DTr} E \in \operatorname{add} E$. Dually, $\operatorname{TrD} E \in \operatorname{add} E$. ■

Remark 3.11. (i) *The condition* $\operatorname{DTr} E \oplus \operatorname{TrD} E \in \operatorname{add} E$ *means that the pairwise non-isomorphic indecomposable direct summands of E are exactly the modules in several* DTr-*obits of finite length.*

(ii) *For a CM-finite algebra, the two conditions in Definition* 3.8 *are in fact equivalent.*

4. Proof of Theorem 1.2

4.1. First we fix some notations.

Put $K^-(\Lambda\text{-mod})$ and $D^-(\Lambda)$ to be respectively the upper bounded homotopy category and the upper bounded derived category of Λ-mod.

Put $K^{-,b}(\mathcal{P}(\Lambda))$ to be the homotopy category of the upper bounded complexes of projective Λ-modules, with only finitely many non-zero cohomology groups. Then there is a canonical triangle-equivalence $D^b(\Lambda) \cong K^{-,b}(\mathcal{P}(\Lambda))$; and Λ is of finite global dimension if and only if there is a canonical triangle-equivalence $D^b(\Lambda) \cong K^b(\mathcal{P}(\Lambda))$.

Put $K^-(\mathcal{GP}(\Lambda))$ to be the upper bounded homotopy category of $\mathcal{GP}(\Lambda)$, and $K^b(\mathcal{GP}(\Lambda))$ the bounded homotopy category of $\mathcal{GP}(\Lambda)$. Denote by $K^{b,ac}(\mathcal{GP}(\Lambda))$ the homotopy category of the bounded acyclic complexes of Gorenstein-projective Λ-modules.

Following 3.3 in [GZ], we define the following triangulated subcategory of $K^-(\mathcal{GP}(\Lambda))$:

$$\begin{aligned} K^{-,gpb}(\mathcal{GP}(\Lambda)) = \{\ G^\bullet \in K^-(\mathcal{GP}(\Lambda)) \mid & \text{ there exists a positive integer } N \text{ such that} \\ & H^{-n}\operatorname{Hom}_\Lambda(G, G^\bullet) = 0,\ \forall\ n > N,\ \forall\ G \in \mathcal{GP}(\Lambda)\}. \end{aligned}$$

Clearly, $K^{-,gpb}(\mathcal{GP}(\Lambda))$ is the Gorenstein version of $K^{-,b}(\mathcal{P}(\Lambda))$, and for any $X^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$, it holds $H^n(X^\bullet) = 0$ for $n << 0$.

4.2. To prove Theorem 1.2, we need some preparations.

Lemma 4.1. *Let $G^\bullet$ be an acyclic complex in $K^{-,gpb}(\mathcal{GP}(\Lambda))$. Then $G^\bullet \in K^{b,ac}(\mathcal{GP}(\Lambda))$.*

Proof. Write $G^\bullet = (G^*, d^*)$. Since $\mathcal{GP}(\Lambda)$ is closed under kernel of epimorphisms, $\operatorname{Im} d^i \in \mathcal{GP}(\Lambda),\ \forall\ i \in \mathbb{Z}$. By definition there exists a positive integer N such that $H^{-n}\operatorname{Hom}(G, G^\bullet) = 0,\ \forall\ n \geq N,\ \forall\ G \in \mathcal{GP}(\Lambda)$. Thus $d^{-(n+1)}$ splits for $n \geq N$, and hence $G^\bullet \in K^{b,ac}(\mathcal{GP}(\Lambda))$. ■

The following observation will play an important role in our argument.

Lemma 4.2. *Let Λ be a CM-finite algebra. Then*

(i) For each $P^\bullet \in K^{-,b}(\mathcal{P}(\Lambda))$, there exists a quasi-isomorphism $P^\bullet \longrightarrow G^\bullet$ with $G^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$.

(ii) For morphisms $f_i^\bullet : P^\bullet \longrightarrow G_i^\bullet$ with $P^\bullet \in K^{-,b}(\mathcal{P}(\Lambda))$ and $G_i^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$, $1 \leq i \leq m$, there exists a quasi-isomorphism $g^\bullet : P^\bullet \longrightarrow G^\bullet$ and chain maps $h_i^\bullet : G^\bullet \longrightarrow G_i^\bullet$ with $G^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$, such that $f_i^\bullet = h_i^\bullet g^\bullet,\ 1 \leq i \leq m$.

Proof. (i) Write $P^\bullet = (P^*, d^*)$. Let N be a positive integer such that $H^{-n}P^\bullet = 0$ for $n \geq N$. By taking right $\mathcal{GP}(\Lambda)$-approximations (this can be done since Λ is CM-finite), we get an exact sequence $\cdots \longrightarrow G^{-N-2} \longrightarrow G^{-N-1} \longrightarrow \operatorname{Ker} d^{-N} \longrightarrow 0$. Combing this with $0 \longrightarrow \operatorname{Ker} d^{-N} \longrightarrow P^{-N} \longrightarrow P^{-N+1} \longrightarrow \cdots$, we get a complex $G^\bullet : \cdots \longrightarrow G^{-N-2} \longrightarrow G^{-N-1} \longrightarrow P^{-N} \longrightarrow P^{-N+1} \longrightarrow \cdots$ in $K^{-,gpb}(\mathcal{GP}(\Lambda))$ with $H^{-n}\operatorname{Hom}(G, G^\bullet) = 0, \forall\ n \geq N,\ \forall\ G \in \mathcal{GP}(\Lambda)$. It is clear that there exists an induced quasi-isomorphism

$$\begin{tikzcd}
P^\bullet = \arrow[d, dashed] & \cdots \arrow[r] & P^{-N-2} \arrow[r] \arrow[d, dashed] & P^{-N-1} \arrow[r] \arrow[d, dashed] & P^{-N} \arrow[r] \arrow[d, equal] & P^{-N+1} \arrow[r] \arrow[d, equal] & \cdots \\
G^\bullet = & \cdots \arrow[r] & G^{-N-2} \arrow[r] & G^{-N-1} \arrow[r] & P^{-N} \arrow[r] & P^{-N+1} \arrow[r] & \cdots .
\end{tikzcd}$$

(ii) Let N be a positive integer such that $H^{-n}\operatorname{Hom}(G, G_i^\bullet) = 0 = H^{-n}P^\bullet$, $\forall\ n \geq N$, $\forall\ G \in \mathcal{GP}(\Lambda)$, $1 \leq i \leq m$. By the proof of (i) we have the following quasi-isomorphism

$$\begin{tikzcd}
P^\bullet = \arrow[d, "g^\bullet"] & \cdots \arrow[r] & P^{-N-2} \arrow[r] \arrow[d] & P^{-N-1} \arrow[r] \arrow[d] & P^{-N} \arrow[r] \arrow[d, equal] & P^{-N+1} \arrow[r] \arrow[d, equal] & \cdots \\
G^\bullet = & \cdots \arrow[r] & G^{-N-2} \arrow[r] & G^{-N-1} \arrow[r] & P^{-N} \arrow[r] & P^{-N+1} \arrow[r] & \cdots
\end{tikzcd}$$

with $G^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$. Since for each i we have $H^{-n}\operatorname{Hom}(G, G_i^\bullet) = 0, \forall\ n \geq N$, $\forall\ G \in \mathcal{GP}(\Lambda)$, by applying $\operatorname{Hom}(G^{-N-1}, -), \operatorname{Hom}(G^{-N-2}, -), \cdots$ to $G_i^\bullet$ we obtain the following chain map $h_i^\bullet$:

$$\begin{tikzcd}
G^\bullet = \arrow[d, dashed, "h_i^\bullet"] & \cdots \arrow[r] & G^{-N-2} \arrow[r] \arrow[d, dashed, "h_i^{-N-2}"] & G^{-N-1} \arrow[r] \arrow[d, dashed, "h_i^{-N-1}"] & P^{-N} \arrow[r] \arrow[d, "f^{-N}"] & P^{-N+1} \arrow[r] \arrow[d, "f^{-N+1}"] & \cdots \\
G_i^\bullet = & \cdots \arrow[r] & G_i^{-N-2} \arrow[r] & G_i^{-N-1} \arrow[r] & G_i^{-N} \arrow[r] & G_i^{-N+1} \arrow[r] & \cdots .
\end{tikzcd}$$

It is easy to see that $f_i^\bullet - h_i^\bullet g^\bullet$ are null homotopy, $1 \leq i \leq m$ (we omit the details). ■

4.3. Consider functor $\rho : K^{-,gpb}(\mathcal{GP}(\Lambda)) \longrightarrow D^-(\Lambda)$, which is the composition of the natural embedding $K^{-,gpb}(\mathcal{GP}(\Lambda)) \hookrightarrow K^-(\Lambda\text{-mod})$ and the standard localization functor $Q : K^-(\Lambda\text{-mod}) \longrightarrow D^-(\Lambda) = K^-(\Lambda\text{-mod})/K^{-,ac}(\Lambda\text{-mod})$, where $K^{-,ac}(\Lambda\text{-mod})$ is the homotopy category of the upper bounded acyclic complexes of Λ-modules.

Since $K^{b,ac}(\mathcal{GP}(\Lambda))$ is a triangulated subcategory of $K^{-,gpb}(\mathcal{GP}(\Lambda))$, we have the Verdier quotient $K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))$, where the corresponding compatible multiplicative system consists of the quasi-isomorphisms in $K^{-,gpb}(\mathcal{GP}(\Lambda))$ with mapping cones being bounded. Since $\rho(K^{b,ac}(\mathcal{GP}(\Lambda))) = 0$, it follows that ρ induces an exact functor $\overline{\rho} : K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)) \longrightarrow D^-(\Lambda)$.

If Λ is CM-finite, then by Lemma 4.2(i) we have $\operatorname{Im}\overline{\rho} = D^b(\Lambda) \cong K^{-,b}(\mathcal{P}(\Lambda))$. So we get an exact functor from $K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))$ to $D^b(\Lambda)$, again denoted by $\overline{\rho}$.

The following proposition is a key step in proving Theorem 1.2.

Proposition 4.3. *Let Λ be a CM-finite algebra. Then the functor*

$$\overline{\rho} : K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)) \longrightarrow D^b(\Lambda)$$

defined above is a triangle-equivalence.

Proof. It remains to prove that $\overline{\rho} : K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)) \longrightarrow D^-(\Lambda)$ is fully faithful. Suppose $G^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$ and $\overline{\rho}(G^\bullet) = 0$. Then $G^\bullet$ is acyclic. By Lemma 4.1 we have $G^\bullet \in K^{b,ac}(\mathcal{GP}(\Lambda))$. That is $\overline{\rho}$ maps non-zero objects to non-zero objects. It is well-known that if an exact functor which sends non-zero objects to non-zero objects is full, then it is faithful. So, it suffices to prove that $\overline{\rho}$ is full.

Let $G_1^\bullet, G_2^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$. Let $\frac{\alpha^\bullet}{s^\bullet} \in \operatorname{Hom}_{D^-(\Lambda)}(\overline{\rho}(G_1^\bullet), \overline{\rho}(G_2^\bullet)) = \operatorname{Hom}_{D^-(\Lambda)}(G_1^\bullet, G_2^\bullet)$, where $s^\bullet : X^\bullet \Longrightarrow G_1^\bullet$ is a quasi-isomorphism with $X^\bullet \in K^-(\Lambda\text{-mod})$, and $\alpha^\bullet : X^\bullet \longrightarrow G_2^\bullet$

is a morphism in $K^-(\Lambda\text{-mod})$. Then there exists a quasi-isomorphism $t^\bullet : P^\bullet \longrightarrow X^\bullet$ with $P^\bullet \in K^-(\mathcal{P}(\Lambda))$, the upper bounded homotopy category of $\mathcal{P}(\Lambda)$. Since $s^\bullet$ and $t^\bullet$ are quasi-isomorphisms and $G_1^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$, it follows that $P^\bullet \in K^{-,b}(\mathcal{P}(\Lambda))$.

Thus we have the following commutative diagram

$$\begin{tikzcd}
 & X^\bullet \arrow[dl, Rightarrow, "s^\bullet"'] \arrow[dr, "\alpha^\bullet"] & \\
G_1^\bullet & P^\bullet \arrow[l, Rightarrow] \arrow[r] \arrow[u, Rightarrow, "t^\bullet"] & G_2^\bullet.
\end{tikzcd}$$

By Lemma 4.2 (ii) we have the following commutative diagram

$$\begin{tikzcd}
G_1^\bullet & P^\bullet \arrow[l, Rightarrow, "s^\bullet t^\bullet"'] \arrow[r, "\alpha^\bullet t^\bullet"] \arrow[d, Rightarrow, "g^\bullet"] & G_2^\bullet \\
 & G^\bullet \arrow[ul, Rightarrow, "l^\bullet"] \arrow[ur, "\beta^\bullet"'] &
\end{tikzcd}$$

where $G^\bullet \in K^{-,gpb}(\mathcal{GP}(\Lambda))$, and $g^\bullet : P^\bullet \longrightarrow G^\bullet$ is a quasi-isomorphism. Note that $l^\bullet$ is also a quasi-isomorphism, and hence the mapping cone $\mathrm{con}(l^\bullet)$ is acyclic. Since $K^{-,gpb}(\mathcal{GP}(\Lambda))$ is a triangulated subcategory of $K^-(\Lambda\text{-mod})$ ([GZ], 3.3), it follows that $\mathrm{con}(l^\bullet) \in K^{-,gpb}(\mathcal{GP}(\Lambda))$. By Lemma 4.1 we have $\mathrm{con}(l^\bullet) \in K^{b,ac}(\mathcal{GP}(\Lambda))$. This proves that $\frac{\beta^\bullet}{l^\bullet} \in \mathrm{Hom}_{K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))}(G_1^\bullet, G_2^\bullet)$ and that $\frac{\alpha^\bullet}{s^\bullet} = \frac{\beta^\bullet}{l^\bullet} = \overline{\rho}(\frac{\beta^\bullet}{l^\bullet})$. This completes the proof. ■

Remark 4.4. (i) *From the proof we see that, the above assertion holds for an arbitrary algebra* Λ *such that* $\mathcal{GP}(\Lambda)$ *is contravariantly finite (i.e., each* Λ*-module has a right* $\mathcal{GP}(\Lambda)$*-approximation, cf. [AR]).*

(ii) *With the same argument, we can prove that there is a triangle-equivalence*

$$K^{-,b}(\mathcal{GP}(\Lambda))/K^{-,ac}(\mathcal{GP}(\Lambda)) \cong D^b(\Lambda)$$

for any Artin algebra (not necessarily CM-finite), where $K^{-,b}(\mathcal{GP}(\Lambda))$ *denotes the homotopy category of the upper bounded complexes of Gorenstein-projective* Λ*-modules, with only finitely many non-zero cohomology groups.*

4.4. We need the following well-known fact.

Lemma 4.5. ([V], *Corollary* 4-3) *Suppose* $\mathcal{D}_1$ *and* $\mathcal{D}_2$ *are triangulated subcategories of triangulated category* $\mathcal{C}$*, and* $\mathcal{D}_1$ *is a subcategory of* $\mathcal{D}_2$*. Then* $\mathcal{D}_2/\mathcal{D}_1$ *is a triangulated subcategory of* $\mathcal{C}/\mathcal{D}_1$*, and there is a triangle-equivalence* $(\mathcal{C}/\mathcal{D}_1)/(\mathcal{D}_2/\mathcal{D}_1) \cong \mathcal{C}/\mathcal{D}_2$.

Denote by $\langle \mathcal{GP}(\Lambda) \rangle$ the triangulated subcategory of $D^b(\Lambda)$ generated by $\mathcal{GP}(\Lambda)$, i.e., the smallest triangulated subcategory of $D^b(\Lambda)$ containing $\mathcal{GP}(\Lambda)$.

Corollary 4.6. *Let Λ be a CM-finite algebra. Then the triangle-equivalence in Proposition* 4.3 $\overline{\rho}: K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)) \longrightarrow D^b(\Lambda)$ *induces a triangle-equivalence*

$$K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)) \longrightarrow \langle \mathcal{GP}(\Lambda)\rangle.$$

Proof. By Lemma 4.5 we have $K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))$ is a triangulated subcategory of $K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))$. Thus $\overline{\rho}(K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)))$ is a triangulated subcategory of $D^b(\Lambda)$. Obviously, it is generated by $\mathcal{GP}(\Lambda)$. ■

Proposition 4.7. *Let Λ be a CM-finite algebra. Then there is a triangle-equivalence*

$$D^b(\Lambda)/\langle \mathcal{GP}(\Lambda)\rangle \cong D^b_{\text{defect}}(\Lambda).$$

Proof. By Buchweitz Theorem ([Buch], Theorem 4.4.1), there is an exact embedding

$$F : \underline{\mathcal{GP}(\Lambda)} \hookrightarrow D^b_{sg}(\Lambda), \quad \text{given by} \quad F(G) = G \tag{1}$$

where the second G is the stalk complex of Gorenstein-projective module G at degree 0.

On other hand, by Lemma 4.5 we know that $\langle \mathcal{GP}(\Lambda)\rangle/K^b(\mathcal{P}(\Lambda))$ is a triangulated subcategory of $D^b_{sg}(\Lambda) = D^b(\Lambda)/K^b(\mathcal{P}(\Lambda))$ and there is a triangle-equivalence

$$D^b(\Lambda)/\langle \mathcal{GP}(\Lambda)\rangle \cong (D^b(\Lambda)/K^b(\mathcal{P}(\Lambda)))/(\langle \mathcal{GP}(\Lambda)\rangle/K^b(\mathcal{P}(\Lambda))). \tag{2}$$

Now we claim

$$\operatorname{Im} F = \langle \mathcal{GP}(\Lambda)\rangle/K^b(\mathcal{P}(\Lambda)). \tag{3}$$

In fact, by Corollary 4.6 in $D^b(\Lambda)$ the objects of $\langle \mathcal{GP}(\Lambda)\rangle$ are isomorphic to bounded complexes with terms in $\mathcal{GP}(\Lambda)$. Note that (1) implies that in $D^b_{sg}(\Lambda)$ the bounded complexes with terms in $\mathcal{GP}(\Lambda)$ are isomorphic to stalk complexes at degree 0 of the modules in $\mathcal{GP}(\Lambda)$. Therefore, in $D^b_{sg}(\Lambda)$ the objects of $\langle \mathcal{GP}(\Lambda)\rangle/K^b(\mathcal{P}(\Lambda))$ are isomorphic to stalk complexes at degree 0 of the modules in $\mathcal{GP}(\Lambda)$, and hence $\langle \mathcal{GP}(\Lambda)\rangle/K^b(\mathcal{P}(\Lambda)) \subseteq \operatorname{Im} F$. On the other hand, it is clear $\operatorname{Im} F \subseteq \langle \mathcal{GP}(\Lambda)\rangle/K^b(\mathcal{P}(\Lambda))$. This proves the claim.

Thus we have

$$\begin{aligned} D^b(\Lambda)/\langle \mathcal{GP}(\Lambda)\rangle &\overset{by\ (2)}{\cong} (D^b(\Lambda)/K^b(\mathcal{P}(\Lambda)))/(\langle \mathcal{GP}(\Lambda)\rangle/K^b(\mathcal{P}(\Lambda))) \\ &\overset{by\ (3)}{=} D^b_{sg}(\Lambda)/\mathrm{Im}F = D^b_{\text{defect}}(\Lambda). \end{aligned}$$

This completes the proof. ■

4.5. **Proof of Theorem 1.2.** By Lemma 4.5, $K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))$ is a triangulated subcategory of $K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))$, and there is a triangle-equivalence

$$\begin{aligned} &K^{-,gpb}(\mathcal{GP}(\Lambda))/K^b(\mathcal{GP}(\Lambda)) \\ &\cong (K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)))/(K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))). \end{aligned} \tag{4}$$

By Corollary 4.6 $\overline{\rho}\,(K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))) = \langle \mathcal{GP}(\Lambda)\rangle$. It follows from Propositions 4.3 and 4.7 that we have the following triangle-equivalences

$$\begin{aligned}&(K^{-,gpb}(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)))/(K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda)))\\ &\cong D^b(\Lambda)/\overline{\rho}\ (K^b(\mathcal{GP}(\Lambda))/K^{b,ac}(\mathcal{GP}(\Lambda))) \cong D^b(\Lambda)/\langle \mathcal{GP}(\Lambda)\rangle \cong D^b_{\text{defect}}(\Lambda).\end{aligned}$$

Together with (4) we get a triangle-equivalence

$$D^b_{\text{defect}}(\Lambda) \cong K^{-,gpb}(\mathcal{GP}(\Lambda))/K^b(\mathcal{GP}(\Lambda)). \tag{5}$$

On the other hand, by the equivalence $\mathcal{P}(\mathbf{AUS}(\Lambda)) \cong \mathcal{GP}(\Lambda)$ of additive categories we get the following triangle-equivalences

$$\begin{aligned}D^b_{sg}(\mathbf{AUS}(\Lambda)) &= D^b(\mathbf{AUS}(\Lambda))/K^b(\mathcal{P}(\mathbf{AUS}(\Lambda)))\\ &\cong K^{-,b}(\mathcal{P}(\mathbf{AUS}(\Lambda)))/K^b(\mathcal{P}(\mathbf{AUS}(\Lambda))\\ &\cong K^{-,gpb}(\mathcal{GP}(\Lambda))/K^b(\mathcal{GP}(\Lambda)).\end{aligned}$$

Together with (5) we have triangle-equivalence $D^b_{\text{defect}}(\Lambda) \cong D^b_{sg}(\mathbf{AUS}(\Lambda))$. This completes the proof. ■

4.6. Final Remark. For an arbitrary algebra Λ such that $\mathcal{GP}(\Lambda)$ is contravariantly finite, by Remark 4.4(i) and by the similar argument we have a triangle-equivalence

$$D^b_{\text{defect}}(\Lambda) \cong K^{-,gpb}(\mathcal{GP}(\Lambda))/K^b(\mathcal{GP}(\Lambda)).$$

In fact, we have the following more general result:

Theorem 4.8. *Suppose Λ and Λ' are two Artin algebras such that $\mathcal{GP}(\Lambda) \cong \mathcal{GP}(\Lambda')$ as additive categories. If $\mathcal{GP}(\Lambda)$ is contravariantly finite, then there is an triangle-equivalence*

$$K^{-,gpb}(\mathcal{GP}(\Lambda'))/K^{b,ac}(\mathcal{GP}(\Lambda')) \longrightarrow D^b(\Lambda').$$

Thus we have triangle-equivalences

$$D^b_{defect}(\Lambda) \cong K^{-,gpb}(\mathcal{GP}(\Lambda))/K^b(\mathcal{GP}(\Lambda)) \cong D^b_{defect}(\Lambda').$$

In particular, Λ is Gorenstein if and only if Λ' is Gorenstein.

Proof. Note that if Λ is Gorenstein then $\mathcal{GP}(\Lambda)$ is contravariantly finite. So the last assertion follows.

We only prove that Λ' satisfies the conclusions (i) and (ii) in Lemma 4.2 (all other arguments are same as the ones for Theorem 1.2). Suppose we have an equivalence $F : \mathcal{GP}(\Lambda) \longrightarrow \mathcal{GP}(\Lambda')$ of categories, with quasi-inverse F^{-1}.

Let $P^\bullet = (P^*, d^*) \in K^{-,b}(\mathcal{P}(\Lambda'))$ with $H^{-n}P^\bullet = 0$ for some integer N and each $n \geq N$. Consider the complex

$$\cdots \longrightarrow F^{-1}(P^{-N-2}) \xrightarrow{F^{-1}(d^{-N-2})} F^{-1}(P^{-N-1}) \xrightarrow{F^{-1}(d^{-N-1})} F^{-1}(P^{-N}) \longrightarrow \cdots.$$

Since $\mathcal{GP}(\Lambda)$ is contravariantly finite, by taking right $\mathcal{GP}(\Lambda)$-approximations we get the following complex

$$G^\bullet : \cdots \longrightarrow G^{-N-3} \xrightarrow{\partial^{-N-3}} G^{-N-2} \xrightarrow{\partial^{-N-2}} F^{-1}(P^{-N-1}) \xrightarrow{F^{-1}(d^{-N-1})} F^{-1}(P^{-N}) \longrightarrow \cdots .$$

By construction we have $H^{-n}\operatorname{Hom}_\Lambda(G, G^\bullet) = 0$ for $G \in \mathcal{GP}(\Lambda)$ and $n \geq N+1$. Since $F : \mathcal{GP}(\Lambda) \longrightarrow \mathcal{GP}(\Lambda')$ is an equivalence, we have the complex isomorphism $\operatorname{Hom}_\Lambda(G, G^\bullet) \cong \operatorname{Hom}_{\Lambda'}(F(G), F(G^\bullet))$ for $G \in \mathcal{GP}(\Lambda)$. It follows that $H^{-n}\operatorname{Hom}_{\Lambda'}(G', F(G^\bullet)) = 0$ for $G' \in \mathcal{GP}(\Lambda')$ and $n \geq N+1$. The rest of the proof is same as the one for Lemma 4.2, and we omit the details. ■

References

[AB] M. Auslander, M. Bridger, Stable module theory, Mem. Amer. Math. Soc. 94., Amer. Math. Soc., Providence, R.I., 1969.

[AR] M. Auslander, I. Reiten, Applications of contravariantly finite subcategories, Adv. Math. 86(1991), 111-152.

[ARS] M. Auslander, I. Reiten, S. O. Smalø, Representation theory of Artin algebras, Cambridge Studies in Adv. Math. 36., Cambridge Univ. Press, 1995.

[B1] A. Beligiannis, The homological theory of contravariantly finite subcategories: Auslander-Buchweitz contexts, Gorenstein categories ans (co-)stabilization, Comm. Algebra 28(10)(2000), 4547-4596.

[B2] A. Beligiannis, Cohen-Macaulay modules, (co)tosion pairs and virtually Gorenstein algebras, J. Algebra 288(1)(2005), 137-211.

[B3] A. Beligiannis, On rings and algebras of finite Cohen-Macaulay type, Adv. Math. 226 (2) (2011), 1973-2019.

[BR] A. Beligiannis, I. Reiten, Homological and homotopical aspects of tosion theories, Mem. Amer. Math. Soc. 188, Amer. Math. Soc., Providence, R.I., 2007.

[BJO] P. A. Bergh, D. A. Jorgensen, S. Oppermann, The Gorenstein defect category, availble in arXiv Math. CT 1202.2876, 2012.

[Buch] R.-O. Buchweitz, Maximal Cohen-Macaulay modules and Tate cohomology over Gorenstein rings, Unpublished manuscript, Hamburg (1987), 155pp.

[BGS] R.-O. Buchweitz, G.-M. Greuel, F.-O. Schreyer, Cohen-Macaulay modules on hypersurface singularities II, Invent. Math. 88(1)(1987), 165-182.

[C1] X. W. Chen, An Auslander-type result for Gorenstein-projective modules, Adv. Math. 218 (2008), 2043-2050.

[C2] X. W. Chen, Algebras with radical square zero are either self-injective or CM-free, Proc. Amer. Math. Soc. 140 (1) (2012), 93-98.

[CPST] L. W. Christensen, G. Piepmeyer, J. Striuli, R. Takahashi, Finite Gorenstein representation type implies simple singularity, Adv. Math. 218(2008), 1012-1026.

[CY] X. W. Chen, Y. Ye, Retractions and Gorenstein homological proerties, availble in arXiv Math. RT 1206.4415, 2012.

[EJ] E. E. Enochs, O. M. G. Jenda, Relative homological algebra, De Gruyter Expo. Math.30 (2000).

[GZ] N. Gao, P. Zhang, Gorenstein derived categories, J. Algebra 323(2010), 2041-2057.

[H1] D. Happel, Triangulated categories in representation theory of finite dimensional algebras, London Math. Soc. Lecture Notes Ser. 119, Cambridge Uni. Press, 1988.

[H2] D. Happel, On Gorenstein algebras, in: Representation theory of finite groups and finite-dimensional algebras, Prog. Math. 95, 389-404, Birkhüser, Basel, 1991.

[Huang] Z. Y. Huang, Proper resolutions and Gorenstein categories, availble in arXiv Math. KT 1203.4110, 2012.

[K] H. Knörrer, Cohen-Macaulay modules on hypersurface singularities I, Invent. Math. 88(1)(1987), 153-164.

[LZ1] Z. W. Li, P. Zhang, Gorenstein algebras of finite Cohen-Macaulay type, Adv. Math. 223 (2010), 728-734.

[LZ2] Z. W. Li, P. Zhang, A construction of Gorenstein-projective modules, J. Algebra 323(2010), 1802-1812.

[R] C. M. Ringel, The Gorenstein-projective modules for the Nakayama algebras, availble in arXiv Math. RT 1207.4856, 2012.

[V] J. L. Verdier, Des catégories dérivées abéliennes, Asterisque 239(1996), xii+253 pp. (1997). With a preface by L. Illusie. Edited and with a note by G. Maltsiniotis.

[SWSW] S. Sather-Wagstaff, T. Sharif, D. White, Stability of Gorenstein categories, J. Math. Soc. 77(2) (2008), 481-502.

[Zhu] S. J. Zhu, Left homotopy theory and Buchweitz's theorem, Master Thesis at SJTU, 2011.